

Clinical Commentary Review

Exposure and Health Effects of Fungi on Humans

Sachin N. Baxi, MD^a, Jay M. Portnoy, MD^b, Désirée Larenas-Linnemann, MD^c, and Wanda Phipatanakul, MD, MS^a; on behalf of the Environmental Allergens Workgroup* Boston, Mass; Kansas City, Mo; and Mexico City, Mexico

Fungi are ubiquitous microorganisms that are present in outdoor and indoor environments. Previous research has found relationships between environmental fungal exposures and human health effects. We reviewed recent articles focused on fungal exposure and dampness as risk factors for respiratory disease development, symptoms, and hypersensitivity. In particular, we reviewed the evidence suggesting that early exposure to dampness or fungi is associated with the development of asthma and increased asthma morbidity. Although outdoor exposure to high concentrations of spores can cause health effects such as asthma attacks in association with thunderstorms, most people appear to be relatively unaffected unless they are sensitized to specific genera. Indoor exposure and dampness, however, appears to be associated with an increased risk of developing asthma in young children and asthma morbidity in individuals who have asthma. These are important issues because they provide a rationale for interventions that might be considered for homes and buildings in which there is increased fungal exposure. In addition to rhinitis and asthma, fungus exposure is associated with a number of other illnesses including allergic bronchopulmonary mycoses, allergic fungal sinusitis, and hypersensitivity pneumonitis. Additional research is necessary to establish causality and evaluate interventions for fungal- and dampness-related health effects. © 2016 American Academy of Allergy, Asthma & Immunology (J Allergy Clin Immunol Pract 2016;■:■-■)

Key words: Fungus; Mold; Health effects; Asthma; Hypersensitivity

Fungi are ubiquitous microorganisms that can be found in all parts of the world. When airborne, fungi take the form of spores, mycelia, and hyphae fragments. Such bioparticulates, when inhaled, are believed to contribute to adverse health effects in individuals who are predisposed to experience disease. Such individuals include those who produce specific IgE to fungal antigens, others with respiratory conditions that are susceptible to irritant effects of exposure, and immunocompromised patients who are susceptible to infections.

Common outdoor fungi include *Alternaria*, *Cladosporium*, *Epicoccum*, as well as ascospores and basidiospores though these fungi are often found indoors because they enter through open doors and windows and can be carried indoors. Fungi more classically associated with indoor water damage or decay include *Penicillium*, *Aspergillus*, *Stachybotrys*, and *Chaetomium*. A review of these diverse fungi can be found in the accompanying article “Taxonomy of allergenic fungi” by Levetin et al.¹

In all epidemiologic studies, exposure to mold and dampness are considered together. Dampness is a generic term that is used to communicate a range of moisture conditions. Although most people can recognize an environment that is damp, there is no consensus description or measurement that defines the term. In this article, *dampness* is defined as sufficient moisture on or in a substrate to support microbial growth. Dampness has often been associated with a combination of the following factors occurring simultaneously: visible water damage or stains, visible mold, and odors from microbial growth.² However, dampness also supports dust mites and sometimes the growth of actinomycetes, which can be associated with other health problems. Thus, it can be challenging to isolate the health effects of fungal exposure in damp environments.³

Another limitation of studies designed to identify health risks of fungal exposure is that total fungal exposure is difficult to measure. There is no criterion standard method to identify and quantify fungus. Methods vary from direct microscopy or culture-based volumetric air sampling to measurement of fungal metabolites such as beta-D-glucan or ergosterol. Fungal diversity can also be evaluated by immunoassays, PCR, and genomic sequencing.⁴ Furthermore, many fungal allergens are broadly cross-reactive.^{5,6} In sensitized individuals, exposure to related species can cause symptoms due to shared epitopes.

In this review, we will address the evidence linking dampness and fungal exposure and adverse health effects. In particular, we will review evidence suggesting that early exposure to dampness or fungi is associated with the development of asthma and that exposure to dampness or fungi in those with asthma increases morbidity. These are important issues because they provide a

^aBoston Children’s Hospital, Boston, Mass

^bChildren’s Mercy Hospital, Kansas City, Mo

^cHospital Médica Sur, Mexico City, Mexico

This study was supported by National Institutes of Health grant numbers K24 AI 106822 and U01 AI 110397.

Conflicts of interest: J. M. Portnoy has received consultancy fees from Church & Dwight. D. Larenas-Linnemann is a board member for the American Academy of Allergy, Asthma & Immunology and Colegio Mexicano de Inmunología Clínica y Alergia; has received consultancy fees from AstraZeneca, Boehringer Ingelheim, MEDA, Chiesi, Pfizer, MSD, Mit Pharma, Novartis, and DBV Technologies; has received lecture fees from AstraZeneca, MEDA, Pfizer, Glenmark, UCB, Novartis, Chiesi, and UCB; has received payment for developing educational presentations from Glenmark; and has received travel support from AstraZeneca, MEDA, Pfizer, Novartis, Sanofi, MSD, and Senosiain. The rest of the authors declare that they have no relevant conflicts of interest.

Received for publication August 18, 2015; revised January 4, 2016; accepted for publication January 13, 2016.

Available online ■■

Corresponding author: Wanda Phipatanakul, MD, MS, Division of Immunology and Allergy, Harvard Medical School, Boston Children’s Hospital, 300 Longwood Ave, Boston, MA 02115. E-mail: wanda.phipatanakul@childrens.harvard.edu.

*The complete list of Environmental Allergens Workgroup members appears in the Acknowledgments.

2213-2198

© 2016 American Academy of Allergy, Asthma & Immunology

<http://dx.doi.org/10.1016/j.jaip.2016.01.008>

Abbreviations used
IOM- Institute of Medicine
OR- odds ratio

rationale for interventions that might be considered for homes and buildings in which there is an increase in occupant fungal exposure. In addition to rhinitis and asthma, exposure to fungi is associated with a number of other illnesses including allergic bronchopulmonary mycoses, allergic fungal sinusitis, and hypersensitivity pneumonitis.

OUTDOOR FUNGAL EXPOSURE

Starting in the 1930s, quantitative measurements of outdoor pollen and spore concentrations have been conducted in the United States.⁷ Quantitation is performed using visual or culture-based air sampling methods, which enumerate the bioaerosol composition of the atmosphere. Volumetric air sampling of fungal spores involves impaction, impingement, or filtration using various instruments. Newer techniques including immunoassays, molecular methods such as PCR, and genomic sequencing are becoming more widely used.⁴

Mean spore concentrations outdoors usually range from 50 spores/m³ during cold, snowy weather to 50,000 or more spores/m³ of air during warm, moist seasons.^{8,9} Because spores can be transported long distances in dust clouds arising in warmer areas,¹⁰ they can be detected in even the most remote regions of the globe.¹¹ Temperature and dew point appear to be important factors that determine the types of spores found in outdoor air.¹² For example, ascospores typically are associated with precipitation, whereas *Alternaria* and *Cladosporium* are associated with dry conditions.

In temperate regions, spores in outdoor air tend to peak in the mid to late summer and decrease with the first hard frost in regions that experience cold winter seasons. Xerophilic spores (eg, *Alternaria*, *Cladosporium*, and *Epicoccum*) tend to peak in the afternoon during periods of low humidity, whereas hydrophilic spores tend to peak during predawn hours when there is high humidity. These include ascospores and basidiospores (mushrooms, puffballs). *Alternaria* is the most prevalent fungus in dry, warm climates.¹³⁻¹⁵

Cladosporium is the most commonly identified outdoor fungus.¹⁶ It is found on dead plants or vegetable matter. *Aspergillus* is often isolated from house dust. It is also found in compost heaps and dead vegetation.¹⁷ *Penicillium* is found in soil, food and grains, and house dust. It grows in water-damaged buildings, wallpaper, and decaying fabrics, often giving a green "mildew" color.

INDOOR FUNGAL EXPOSURE

Indoor fungal taxa are likely to vary depending on building construction and climate. The most common taxa are a combination of fungi classically associated with dampness as well as outdoor fungi that enter through openings in the building such as doors and windows.

In a study of 23 buildings, water leakage through roofs, dampness, and defective plumbing were the main reasons for damage leading to fungal contamination.¹⁸ In a study of damp buildings, the genera most frequently encountered indoors included *Penicillium* (68%), *Aspergillus* (56%), *Chaetomium* (an

ascospore) (22%), *Ulocladium* (which is related to *Alternaria*) (21%), *Stachybotrys* (19%), and *Cladosporium* (15%).¹⁸ One study of 21 offices in 4 office buildings showed that concentrations of dustborne fungi positively related to carbon dioxide and were highest at temperatures between 20°C and 22.5°C. In addition, fungal concentrations were highest in September and lowest in March.¹⁹

In a survey of 190 homes in Paris, the most common indoor isolates were *Cladosporium* and *Penicillium* species. *Aspergillus* was recovered in 60% of homes and *Alternaria* in less than 20%. The best predictors for indoor fungal concentrations were their outdoor concentrations when windows were open and the overall dampness in the house when windows were closed.²⁰ In recent studies exploring the fungal species spectrum in 173 homes in the Midwest region of the United States, *Cladosporium*, *Penicillium*, *Aspergillus*, *Basidiospores*, *Epicoccum*, and *Pithomyces* were more frequently present and in higher concentrations in homes with a child who has asthma than in homes without a child with asthma even after adjusting outdoor spore concentration.²¹

Building products most vulnerable to mold attacks include organic materials containing cellulose, wood, jute, wallpaper, drywall, and cardboard.¹⁸ *Penicillium* can use various nutrients such as decaying vegetation with relatively little dampness for short periods of time, whereas *Stachybotrys* requires sustained wetness and easily digestible cellulose such as drywall or paper. Despite these variations in growth requirements, there is a surprising overall consistency in the types of indoor spores.

One study demonstrated that rankings by prevalence and abundance of the types of airborne and dustborne fungi did not differ from winter to summer, nor did their indoor to outdoor ratios differ.²² During the winter when infiltration of spores from outdoors was minimal, mean indoor levels of airborne spores in one study ranged from less than 10 spores/m³ to more than 20,000 spores/m³.²³

INDOOR FUNGAL EXPOSURE AND THE RISK OF DEVELOPING ASTHMA

Studies of the possible association between early fungal exposure and subsequent development of asthma or rhinitis have been performed. Such studies are necessarily observational because it is not feasible to randomly assign children to live in environments with various amounts of fungal exposure and to then monitor their health over time. Although observational studies provide information about associations, they cannot be used to infer causality because other factors that correlate with fungal exposure may also influence any observed health effects.

A systematic review by Mendell et al² evaluated various types of fungal exposure and the risk of developing asthma. The authors identified 17 studies of fungal exposure and subsequent asthma (8 that had been included in an earlier Institute of Medicine [IOM] report²⁴ and 9 new studies) that met their inclusion criteria. Of these studies, 6 were prospective, 8 were retrospective, and 3 were cross-sectional. The odds ratios (ORs) for the development of asthma in these studies ranged from 0.63 to 7.08 in both retrospective and prospective studies; however, only the cross-sectional studies included OR ranges that were consistently greater than 1. They concluded that evidence showed indoor dampness or mold to be associated with asthma development. The limitation of this review is that the studies were not quantitatively

evaluated or combined into a meta-analysis, so it is difficult to draw conclusions from the combined results of the various studies.

The authors did call attention to the Pekkanen et al²⁵ study, which was a case-control study of 121 children 12 to 84 months old with recent onset of wheezing and 241 controls. Homes were inspected by engineers for evidence of moisture, visible mold, moldy odor, and water damage. Although dampness or mold in the main living area of houses was found to be associated with asthma onset, most of the correlations for other interior spaces were not significant. A methodological criticism of this study is that the home inspections were done after asthma was diagnosed; thus, it is difficult to differentiate between asthma development versus worsening. This demonstrates the limitations of observational studies of fungal exposure and asthma onset.

A number of prospective cohort studies of fungal exposure and subsequent development of asthma were recently evaluated by Quansah et al²⁶ in a systematic review. In this review, the authors identified 16 studies that evaluated the relationship between dampness or mold exposure and the development of asthma. Of these, 8 evaluated water damage, 9 evaluated dampness, 12 evaluated visible mold, and 8 looked at moldy odor. There were no studies meeting inclusion criteria that evaluated fungal exposure and development of rhinitis. Summary effects (a combination of ORs and incidence rate ratios) of these studies did not show a significant relationship between water damage and subsequent development of asthma. However, significant summary effects were found for dampness (1.33; CI, 1.12-1.56), visible mold (1.29; CI, 1.04-1.60), and moldy odor (1.73; CI, 1.19-2.50). The authors suggested that each of these types of exposure can be placed into a sequence of occurrence, with water damage leading to dampness, dampness leading to visible mold, and visible mold leading to moldy odor, though the clinical relevance of this postulated sequence is unclear unless it suggests a way to prevent progression to the next step.

Karvonen et al²⁷ recently reported the results of a prospective birth cohort study evaluating the association between moisture damage and visible mold exposure during infancy and the development of asthma up to the age of 6 years. A total of 442 children were followed. A trained civil engineer conducted a home inspection when the child was 5 months old. Moisture damage was classified as no damage, minor damage, or major damage, and the area was quantified and the presence of mold odor or visible mold was noted. Questionnaires were used at 12, 18, and 24 months and then annually to obtain information about respiratory symptoms and physician-diagnosed asthma. During the 6-year follow-up, 65 children developed asthma and 35 children had persistent asthma at 6 years. The strongest associations, with dose-related responses, were found between persistent asthma and moisture damage, with visible mold in the child's bedroom (OR, 4.82; CI, 1.29-18.02) and in the living room (OR, 7.51; CI, 1.49-37.83). The association was stronger if asthma was diagnosed at age 2 years or younger and in atopic children. The results indicate that exposure to moisture damage and mold at an early age is associated with asthma ever and persistent asthma.

The conclusion to take from these studies is that early exposure to environments with dampness, visible mold, and moldy odor is associated with subsequent development of asthma. What is still not known is how much exposure is required (amount and duration), whether there is a dose response, whether specific genera of fungi are responsible for the effect, and whether

interventions to reduce exposure would prevent the development of asthma. We also do not know whether certain populations are more sensitive to exposure. These are areas that require more study. In the meantime, it seems to the authors that it would be prudent for young children to avoid excessive moisture damage and fungal exposure as much as is feasible until further evidence becomes available.

INDOOR FUNGAL EXPOSURE AND CURRENT ASTHMA

A great deal of study has gone into the potential adverse health effects of indoor fungal exposure. Such studies are complicated by the fact that measurement of personal exposure in terms of amount and duration is difficult, many different fungal species contribute to the exposure, and sensitivity to the exposure is likely to vary depending on the person who is exposed. In addition, the only way to infer that exposure to fungi is a cause of observed health effects is to prospectively, in a randomized controlled manner, expose patients to fungi or to reduce fungal exposure in symptomatic patients to document improvement. Given the difficulty of performing such investigations, virtually all studies have been observational.

The systematic review by Mendell et al² evaluated in detail various types of fungal exposure and the risk of developing specific asthma symptoms. The authors identified 103 studies that included 16 health outcomes. Types of exposure reported in these studies included visible water damage, dampness, leaks, flooding, visible condensation, visible mold, and moldy odor. Fewer studies evaluated more quantitative measures of fungal exposure such as culturable or microscopically identified fungi, ergosterol, extracellular polysaccharides, (1 → 3)- β -d-glucans, endotoxin, or lipopolysaccharides. They concluded that evidence from epidemiologic studies and meta-analyses showed indoor dampness or mold to be associated consistently with increased asthma exacerbations, dyspnea, wheeze, and cough, and evidence strongly suggested causation of asthma exacerbations in children.

An IOM meta-analysis of the 33 studies evaluated health outcomes such as upper respiratory tract symptoms, wheeze, cough, and asthma diagnosis in relation to visible mold, dampness, and airborne spores.²⁴ These showed evidence of an association between the presence of fungal growth and dampness and upper respiratory tract symptoms, cough, wheeze, and asthma. Fisk et al²⁸ reported results of quantitative meta-analyses of these studies from the IOM report plus other related studies and concluded that building dampness and fungal growth are associated with increases of 30% to 50% in cough, wheeze, and asthma.

A recent Cochrane meta-analysis by Sauni et al²⁹ evaluated the evidence from controlled clinical studies and trials on the role of interventions geared toward reducing mold exposure and a reduction in asthma symptoms. They found that repairing mold-damaged houses, schools, and offices decreased asthma-related symptoms and respiratory infections compared with no intervention in adults but not in children and suggested that there is a need for better randomized controlled trials to provide more evidence. By inference, this meta-analysis provides evidence that interventions that reduce mold exposure can possibly improve asthma symptoms and suggests a linkage between health effects and exposure in adults but further studies in children are needed.²⁹ A detailed review of interventions is discussed elsewhere in this issue.

A study of 4600 children from the Harvard Six Cities Study that evaluated 10-year-old children with a questionnaire found that wheeze and cough were significantly associated with the presence of fungi and dampness.³⁰ The Health Canada study included nearly 13,000 children and 15,000 adults from the interior of British Columbia to Nova Scotia³¹ who lived in single-family detached houses (81%), small apartment buildings (13%), or single-family attached homes (6%). Visible fungi were reported in 32% of homes, flooding in 24%, and moisture in 14%. For children, bronchitis, cough, and increased wheeze were found to be more frequent in homes with reported fungi or dampness.³¹ For adults, the presence of home dampness and visible fungi was associated with increased prevalence of lower respiratory tract symptoms regardless of other factors or the presence of allergies (OR, 1.62; CI, 1.48-1.78).³²

A recent review by Kanchongkittiphon et al³³ provided an update to the 2000 review of environmental exposures by the IOM. The authors identified 6 studies that provided updated evidence on associations between fungal exposure and asthma morbidity. All the studies were done in children: 5 were prospective, and 1 was cross-sectional. The 2 most recent studies in the review were prospective studies that found that in fungus-sensitized children, indoor *Penicillium* was associated with increased asthma exacerbations.^{34,35} The study by Gent et al³⁴ found that children sensitized and exposed to low levels of *Penicillium* had increased morbidity. Pongracic et al³⁵ found that there was a significant increase in maximum symptom days per 2 weeks in fungus-sensitized inner-city children with asthma exposed to indoor *Penicillium* (1.19 maximum symptom days per 2 weeks; $P = .03$). A prospective study by Bundy et al³⁶ found indoor *Penicillium* to be associated with increased peak expiratory flow variability greater than 18.5% (OR, 2.4; CI, 1.2-4.8). The authors of the review concluded that there was limited evidence of an association between indoor culturable *Penicillium* exposure and exacerbations in children with asthma.

ALLERGIC SENSITIZATION TO FUNGI AND HEALTH EFFECTS

Some of the above studies found an association between exposure to fungi and the risk of respiratory symptoms regardless of sensitization (ie, the presence of fungus-specific IgE). However, sensitization to fungi may increase the risk of morbidity as noted in the studies by Gent et al³⁴ and Pongracic et al³⁵ as described above though most of the evidence is related to outdoor fungal sensitivities. In one study, patients admitted to an intensive care unit for asthma were more likely to have a positive skin test to fungi but not to grass mix, cat dander, or house dust mite.³⁷ In adults with asthma, skin prick test positivity to 5 fungi (*Aspergillus*, *Alternaria*, *Cladosporium*, *Penicillium*, and *Candida*) and 4 other common allergens was explored in relation to asthma severity as measured by the number of hospital admissions. A total of 76% of patients with multiple admissions had at least 1 positive fungal skin test result compared with 16% to 19% of other patients with asthma, and multiple fungal skin test positivity and larger skin test reactions to fungal allergen were also correlated to multiple hospital admissions.³⁸ Sensitivity to *Aspergillus fumigatus* has been associated with severe persistent asthma in adults.³⁹

Alternaria sensitivity has been found to be a risk factor for severe asthma attacks and epidemic asthma. In one study,

Chicago asthma deaths were more than 2 times higher on days when there were 1000 *Alternaria* spores/m³ of air.¹⁵ In another study, increased exposure to *Alternaria* was associated with sudden respiratory arrest in 11 *Alternaria*-sensitive patients with asthma at Mayo Clinic.¹⁴ Another study found that outdoor increases in exposure to *Cladosporium* and *Epicoccum* also were correlated with reductions in peak flow and increased cough in children with asthma, independent of total particle counts and air pollution.⁴⁰

Fungal spores and bioaerosols also are suspected to be associated with thunderstorm-induced asthma epidemics.⁴¹⁻⁴⁵ For example, in the United Kingdom, an epidemic of 26 asthma admissions was associated with a thunderstorm that contained increased levels of *Alternaria*, *Cladosporium*, and *Didymella* species. Most of the patients had *Alternaria*-specific IgE, suggesting that allergy to *Alternaria* may have been the cause.¹³

Tables I and II highlight recent studies that summarize evidence of the relation of fungal exposure and asthma and allergic symptoms.

OTHER HYPERSENSITIVITY DISORDERS

Several other well-defined human diseases are known to be caused or triggered by exposure to fungi or their metabolites. These include allergic bronchopulmonary mycosis, allergic fungal sinusitis, and hypersensitivity pneumonitis.⁶¹ Although exposure to fungi clearly can worsen each of these conditions, complete fungal avoidance has not been shown to result in clinical improvement. The goal of environmental control for these disorders therefore is to avoid making them worse.

Allergic bronchopulmonary mycosis is an inflammatory lung disease that is characterized by a constellation of criteria that include asthma, fleeting pulmonary opacities, proximal bronchiectasis, eosinophilia, and elevated total IgE level in addition to elevated specific IgE level and precipitating IgG to certain fungi. It often occurs in individuals with asthma or cystic fibrosis.⁶² The most common fungus associated with this disorder is *Aspergillus*. More recently, it has been described with other genera of fungi, leading to the generic term "bronchopulmonary mycosis."^{63,64} This disorder is exacerbated by exposure to fungal allergens to which the patient is sensitive, making avoidance an important component of management. Conventional wisdom is that exposure to the offending fungus genus is associated with increased disease severity; however, there have not been any controlled trials of fungal avoidance in this disorder. Extensive reviews of this condition have been published recently.^{65,66}

Allergic fungal sinusitis is a distinct noninvasive fungal sinusitis that is underdiagnosed, with the incidence varying by region of the country.⁶⁷ The offending organisms typically include taxa such as *Alternaria*, *Epicoccum*, *Ulocladium*, *Botrytis*, and *Bipolaris*. The presence of nasal polyps has been reported in conjunction with allergic fungal sinusitis and sensitivity to fungi as evidenced by elevated specific IgE levels.⁶⁷ This condition represents an intense allergic response against fungi, giving rise to the formation of allergic (eosinophilic) mucin, mucostasis, and sinus opacification.⁶⁸ Several potential deficits in the innate and acquired immunity of patients with fungal sinusitis appear to alter host ability to respond to fungi. It is also possible that fungi have a disease-modifying role in the development of this

TABLE I. Recent studies evaluating the association of culturable fungi with asthma exacerbations or severity

Source	Study design	Measured exposures	Findings
Children			
Dannemiller et al ⁴⁶	Birth cohort	Fungal taxa and diversity	They found that a lower fungal diversity demonstrated significant associations with the development of childhood asthma, though they did not report about fungal quantity.
Gent et al ³⁴	Prospective (over 1 mo)	Culturable (1-min samples) airborne fungi (<i>Penicillium</i> , <i>Cladosporium</i>) in living room	Among specifically sensitized children with asthma, relative to unexposed or exposed nonsensitized, significant positive association of any <i>Penicillium</i> exposure with doubled levels of increased asthma severity score, wheeze, and persistent cough. No associations seen with <i>Cladosporium</i> .
Pongracic et al ³⁵	Prospective cohort	Indoor and outdoor culturable (1-min samples) airborne fungi levels measured at baseline and throughout the 2-y study	Among children with asthma with any fungal sensitization, total indoor fungi, the 4 most common fungi combined (<i>Alternaria</i> , <i>Aspergillus</i> , <i>Cladosporium</i> , and <i>Penicillium</i>), and indoor <i>Penicillium</i> were associated with increased severe exacerbations after controlling for outdoor exposure; <i>Penicillium</i> exposure was associated with increased symptoms among children who were fungus-sensitized but not to <i>Penicillium</i> . Among species, only <i>Penicillium</i> exposure demonstrated significant effects among nonspecifically sensitized. The sum of 4 most common indoor fungi and indoor <i>Penicillium</i> was associated with increased symptom days. In conclusion, outdoor and indoor fungi, especially <i>Penicillium</i> , worsen asthma morbidity in inner-city children. Indoor <i>Penicillium</i> uniquely affected both symptoms and UVs.
Wu et al ⁴⁷	Prospective cohort study (outcomes every 4 mo over 4 y)	Total cultural fungi/in vacuumed dust from 5 home locations at initial visit, and genetic polymorphisms	For high values of total culturable fungi in house dust, although not directly related to outcomes of urgent care visits, lung function, IgE, or eosinophils, relationships with urgent care visits were significantly modified by 3 SNPs of chitinase in unadjusted analyses. (As estimated from a figure, the association of high vs low fungi with severe exacerbations [$\sim 30\%$ vs 0%] in those with no or 1 copy of SNP rs2486953 approximately doubled with 2 copies.) Thus, reduced enzymatic breakdown of fungal chitin may increase susceptibility to effects of chitin.
Bundy et al ³⁶	Prospective (over 2 wk)	Culturable (1-min samples) airborne <i>Penicillium</i> and <i>Cladosporium</i> in main living area of home	Any measured culturable airborne <i>Penicillium</i> in the main living area was significantly associated in children with asthma (unknown sensitization, but adjusted for maternal-reported atopy) with more than doubled PEFV over the next 2 wk; exposure to any <i>Alternaria</i> was associated nonsignificantly with twice the likelihood to have the highest PEFV. No associations were found with total mold, <i>Cladosporium</i> , or <i>Aspergillus</i> . Analyses atopy-adjusted.
Inal et al ⁴⁸	Prospective (monthly over 1 y)	Culturable (time unspecified) airborne fungi in living room and bedroom	In children with asthma (17 of 19) and/or rhinitis, sensitized only to molds, in <i>unadjusted analyses in 19 children</i> , total indoor molds, <i>Cladosporium</i> , <i>Alternaria</i> , <i>Penicillium</i> , or <i>Aspergillus</i> had significant correlations with daily asthma symptom score, morning PEF, or evening PEF.
Turyk et al ⁴⁹	Cross-sectional	Culturable (time unspecified) airborne fungi in kitchen and bedroom	Bedroom <i>Penicillium</i> , in models adjusted for home dampness, had a significantly positive (dose-response) association with frequent asthma symptoms in children with asthma (unknown atopic status), and a nearly significant positive association with the number of asthma symptoms. No associations seen for kitchen <i>Penicillium</i> .

PEF, Peak expiratory flow; PEFR, peak expiratory flow rate; PEFV, peak expiratory flow variability; SNPs, single nucleotide polymorphisms; UVs, unscheduled medical visits.

condition.⁶⁹ Several reviews of allergic fungal sinusitis have been published recently.⁷⁰⁻⁷²

Hypersensitivity pneumonitis is an inflammatory lung disease that is caused by an exaggerated immune response to the inhalation of a large variety of organic particles. The most frequent causal antigens are bird proteins and bacteria such as *thermophilic*

actinomycetes. Fungi also have been implicated in both occupational and nonoccupational outbreaks.⁷³ When fungi are involved, it is characterized by the presence of precipitating IgG antibodies directed at the fungus.⁷⁴ The clinical course of the disease is variable, and its diagnosis is difficult because no specific test or biomarker provides a consistent diagnosis. The

TABLE II. Recent studies evaluating dampness, visible mold, or dampness-related agents with asthma exacerbations

Source	Study design	Measured exposures	Findings
Children			
Karvonen et al ²⁷	Birth cohort	Moisture damage	They showed that moisture damage and mold exposure at an early age in the child's main living areas were associated with asthma development: Associations with asthma ever were strongest for visible mold in the child's bedroom (OR of 4.82 [CI, 1.29-18.02]) and in the living room (OR of 7.51 [CI, 1.49-37.83]).
Tiesler et al ⁵⁰	Birth cohort	Visible mold	German-based population study found that visible mold or dampness at home negatively influenced sleep in children. Results were significant for any sleep problem, problems sleeping through the night, and short-term sleep.
Weinmayer et al ⁵¹	Cross-sectional	Damp housing	International Study of Asthma and Allergies in Childhood phase 2, cross-sectional study of 8-12-y-olds in 20 countries evaluating the influence of damp housing conditions on upper and lower respiratory tract symptoms. Damp housing was associated with wheezing in the past year (OR 1.58 [CI, 1.4-1.79]), also significant for occurrence of severe wheeze, speech-limiting wheeze, and ≥ 4 attacks per week. Exposure to dampness, both at present and earlier in life, was associated with wheeze occurrence.
Hagmolen of Ten Have et al ⁵²	Prospective	Parentally reported damp stains or mold growth, in living room or bedroom, in last 2 y	Damp stains or mold growth was associated with significant increases in severe airway hyperresponsiveness, more days with respiratory symptoms, and greater PEF variability.
Kercsmar et al ⁵³	Randomized controlled intervention	Remediation of root causes of home moisture and mold, removal of water-damaged building materials, and cleaning	In the period after remediation, relative to the control group, visible mold scores were significantly more reduced (-2.6 vs -1.4), and measured mold indices were nonsignificantly reduced (-0.41 vs 0.33); in the remediation group compared with the control group, maximum symptom days were significantly reduced and subjects having 1+ acute care visits were reduced by 64% in the as-randomized analysis, and by 86% in the as-treated analysis. About one-third of the subjects were sensitive to fungi.
Bernstein et al ⁵⁴	Controlled intervention	Intervention: 2-wk ultraviolet radiation to reduce microbial exposures	In children sensitized to fungi, ultraviolet radiation was associated with a significant reduction in PEF variability and nonsignificant reduction in FEV ₁ ; significant reductions in severity scores for shortness of breath and chest tightness, and in number of days of shortness of breath and chest tightness, as well as in amount of medication use; nonsignificant reductions in all other disease severity measures.
Bonner et al ⁵⁵	Cross-sectional	Presence of any moisture or mildew	Presence of moisture or mildew at home associated with 3.31 times more hospitalization visits for breathing-related problems, 3.25 times more frequent wheezing episodes, and expected 2.19 times greater frequency of night symptoms.
Teach et al ⁵⁶	Cross-sectional	Mold or dampness in the home in the previous month	Visible dampness or mold in the home in the previous month, in children with asthma of unknown sensitization status, was not associated with unscheduled visits above the median, persistent asthma symptoms, or quality-of-life scores below the median.
Venn et al ⁵⁷	Prospective	Observed mold and measured wall moisture	In children with persistent wheezing across a 3-y interval, there was a <i>dose-related</i> increase in wheezing with increasing measured wall dampness, significantly more in atopic cases. For nighttime symptoms and bedroom dampness, the OR was 2.51 (1.36-4.64) per increasing category, with OR = 7.0 for the highest category; for daytime symptoms and living room dampness, the OR was 1.86 (1.02-3.42) per increasing category. Visible mold was not significantly associated with either symptoms, although it was significantly associated with wheezing illness.

(continued)

TABLE II. (Continued)

Source	Study design	Measured exposures	Findings
Adults			
Jaakkola et al ⁵⁸	Meta-analysis	Mold odor or visible mold	Meta-analysis of 31 studies looking at the relationship of mold odor or visible mold and rhinitis. There was a significant OR for mold odor and rhinitis (OR, 2.18) and allergic rhinitis (OR, 1.87). Also significant was the presence of mold and rhinitis (1.82) and allergic rhinitis (1.51).
Wen et al ⁵⁹	Cross-sectional	Visible mold from interview	Prevalence of asthma attacks among those exposed to indoor mold was roughly twice that in those not exposed, in either obese or nonobese subjects.
Williamson et al ⁶⁰	Prospective	Inspector-assessed visible mold, and measured wall moisture as total dampness or worst dampness	In those in whom asthma was diagnosed, asthma severity had a significant positive <i>dose-related</i> association with measured total dampness and with visible mold score; measured airflow obstruction was significantly greater with higher measured dampness.

PEF, Peak expiratory flow; PEFR, peak expiratory flow rate.

histopathology usually consists of a granulomatous interstitial bronchiolocentric pneumonitis characterized by the presence of poorly formed granulomas and a prominent interstitial infiltrate composed of lymphocytes, plasma cells, and macrophages.⁷⁵

In a German study, 23 children with confirmed hypersensitivity pneumonitis were identified in 2005-2006 and fungal sensitivity was second only to bird sensitivity as the suspected sensitizing agent.⁷⁶ There is also a well-documented pediatric case in which home exposure to *Aspergillus* was demonstrated.⁷⁷ Similar cases have been reported in adults for other fungi including taxa such as *Cladosporium*.⁷⁸ Morbidity clearly is increased if exposure to the offending substance persists. What is less clear is whether having the patient avoid exposure to the offending agent can reduce morbidity. Several reviews of hypersensitivity pneumonitis have been published recently.^{79,80}

SUMMARY

Fungi are ubiquitous microorganism that are present in both outdoor and indoor air. Although outdoor exposure to high concentrations of spores can cause health effects such as asthma attacks in association with thunderstorms, most people appear to be relatively unaffected unless they are sensitized to specific genera. Indoor exposure and dampness, however, appear to be associated with an increased risk of developing asthma in young children, and with asthma morbidity in individuals who have asthma. Reduced indoor exposure using various interventions primarily aimed at reducing moisture, killing fungi, and removing contaminated materials has been shown to decrease this risk of morbidity.

What is not known is how much exposure is necessary to cause a particular health effect and whether certain species are more likely to cause such effects. This review summarizes recent studies pointing toward health effects related to mold. Future studies determining the most accurate method to identify and quantify fungi may help answer the remaining questions.

Acknowledgments

Environmental Allergens Workgroup members: Charles Barnes, PhD, Center for Environmental Health, the Children's Mercy Hospitals & Clinics, Kansas City, Mo; Sachin Baxi, MD, Immunology, Boston Children's Hospital, Boston, Mass; Carl Grimes, CIEC, Healthy Habitats LLC, Denver, Colo; W. Elliott

Horner, PhD, UL Environment, Marietta, Ga; Kevin Kennedy, MPH, CIEC, Center for Environmental Health, Children's Mercy Hospital, Kansas City, Mo; Desiree Larenas-Linnemann, MD, Allergy Faculty, Hospital Medica Sur, Mexico City, Mexico; Estelle Levetin, PhD, Faculty of Biological Science, The University of Tulsa, Tulsa, Okla; J. David Miller, PhD, NSERC Industrial Research Chair, Carlton University, Ottawa, Ontario, Canada; Wanda Phipatanakul, MD, MS, Division of Allergy and Immunology, Harvard Medical School, Boston Children's Hospital, Boston, Mass; Jay M. Portnoy, MD, Division of Allergy, Asthma & Immunology, Children's Mercy Hospital and University of Missouri-Kansas City School of Medicine, Kansas City, Mo; James Scott, PhD, ARMCCM, Division of Occupational & Environmental Health, Dalla Lana School of Public Health, University of Toronto, Toronto, Ontario, Canada; and P. Brock Williams, PhD, Division of Allergy, Asthma & Immunology, University of Missouri-Kansas City School of Medicine and the Children's Mercy Hospitals & Clinics, Kansas City, Mo.

REFERENCES

- Levetin E, Horner WE, Scott JA, et al. Environmental Allergens Workgroup. Taxonomy of allergenic fungi. *J Allergy Clin Immunol Pract* 2016.
- Mendell MJ, Mirer AG, Cheung K, Tong M, Douwes J. Respiratory and allergic health effects of dampness, mold, and dampness-related agents: a review of the epidemiologic evidence. *Environ Health Perspect* 2011;119:748-56.
- WHO guidelines for indoor air quality: dampness and mould. Copenhagen, Denmark: World Health Organization; 2009.
- Levetin E. Methods for aeroallergen sampling. *Curr Allergy Asthma Rep* 2004; 4:376-83.
- Chou H, Tam MF, Chiang CH, Chou CT, Tai HY, Shen HD. Transaldolases are novel and immunoglobulin E cross-reacting fungal allergens. *Clin Exp Allergy* 2011;41:739-49.
- Soeria-Atmadja D, Onell A, Borga A. IgE sensitization to fungi mirrors fungal phylogenetic systematics. *J Allergy Clin Immunol* 2010;125:1379-1386.e1.
- Durham OC. Pollen and hay fever. *Science* 1938;87:167-8.
- Pashley CH, Fairs A, Free RC, Wardlaw AJ. DNA analysis of outdoor air reveals a high degree of fungal diversity, temporal variability, and genera not seen by spore morphology. *Fungal Biol* 2012;116:214-24.
- Barnes C, Schreiber K, Pacheco F, Landuyt J, Hu F, Portnoy J. Comparison of outdoor allergenic particles and allergen levels. *Ann Allergy Asthma Immunol* 2000;84:47-54.
- Darwin C. An account of the fine dust which often falls on vessels in the Atlantic Ocean. *R J Geol Soc* 1846;2:26-30.
- Meier F, Lindbergh C. Collecting micro-organisms from the Arctic atmosphere: with field notes and material. *Sci Monthly* 1935;40:4-20.

12. Troutt C, Levetin E. Correlation of spring spore concentrations and meteorological conditions in Tulsa, Oklahoma. *Int J Biometeorol* 2001;45:64-74.
13. Pulimood TB, Corden JM, Bryden C, Sharples L, Nasser SM. Epidemic asthma and the role of the fungal mold *Alternaria alternata*. *J Allergy Clin Immunol* 2007;120:610-7.
14. O'Hollaren MT, Yunginger JW, Offord KP, Somers MJ, O'Connell EJ, Ballard DJ, et al. Exposure to an aeroallergen as a possible precipitating factor in respiratory arrest in young patients with asthma. *N Engl J Med* 1991;324:359-63.
15. Targonski PV, Persky VW, Ramekrishnan V. Effect of environmental molds on risk of death from asthma during the pollen season. *J Allergy Clin Immunol* 1995;95:955-61.
16. Bensch K, Braun U, Groenewald JZ, Crous PW. The genus *Cladosporium*. *Stud Mycol* 2012;72:1-401.
17. Kumar R, Verma D, Singh BL, Kumar U, Shweta. Composting of sugar-cane waste by-products through treatment with microorganisms and subsequent vermicomposting. *Bioresour Technol* 2010;101:6707-11.
18. Gravesen S, Nielsen PA, Iversen R, Nielsen KF. Microfungal contamination of damp buildings—examples of risk constructions and risk materials. *Environ Health Perspect* 1999;107:505-8.
19. Chao HJ, Milton DK, Schwartz J, Burge HA. Dustborne fungi in large office buildings. *Mycopathologia* 2002;154:93-106.
20. Dassonville C, Demattei C, Detaint B, Barral S, Bex-Capelle V, Momas I. Assessment and predictors determination of indoor airborne fungal concentrations in Paris newborn babies' homes. *Environ Res* 2008;108:80-5.
21. Meng J, Barnes CS, Rosenwasser LJ. Children's Mercy Center for Environmental Health. Identity of the fungal species present in the homes of asthmatic children. *Clin Exp Allergy* 2012;42:1448-58.
22. Horner WE, Worthan AG, Morey PR. Air- and dustborne mycoflora in houses free of water damage and fungal growth. *Appl Environ Microbiol* 2004;70:6394-400.
23. Solomon WR. A volumetric study of winter fungus prevalence in the air of midwestern homes. *J Allergy Clin Immunol* 1976;57:46-55.
24. Institute of Medicine. Damp indoor spaces and health. Washington, DC: National Academy of Sciences, Board on Health Promotion and Disease Prevention, National Academies Press; 2004.
25. Pekkanen J, Hyvarinen A, Haverinen-Shaughnessy U, Korppi M, Putus T, Nevalainen A. Moisture damage and childhood asthma: a population-based incident case-control study. *Eur Respir J* 2007;29:509-15.
26. Quansah R, Jaakkola MS, Hugg TT, Heikkinen SA, Jaakkola JJ. Residential dampness and molds and the risk of developing asthma: a systematic review and meta-analysis. *PLoS One* 2012;7:e47526.
27. Karvonen AM, Hyvarinen A, Korppi M, Haverinen-Shaughnessy U, Renz H, Pfefferle PI, et al. Moisture damage and asthma: a birth cohort study. *Pediatrics* 2015;135:e598-606.
28. Fisk WJ, Lei-Gomez Q, Mendell MJ. Meta-analyses of the associations of respiratory health effects with dampness and mold in homes. *Indoor Air* 2007;17:284-96.
29. Sauni R, Verbeek JH, Uitti J, Jauhiainen M, Kreiss K, Sigsgaard T. Remediating buildings damaged by dampness and mould for preventing or reducing respiratory tract symptoms, infections and asthma. *Cochrane Database Syst Rev* 2015;(2):CD007897.
30. Brunekreef B, Dockery DW, Speizer FE, Ware JH, Spengler JD, Ferris BG. Home dampness and respiratory morbidity in children. *Am Rev Respir Dis* 1989;140:1363-7.
31. Dales RE, Zwanenburg H, Burnett R, Franklin CA. Respiratory health effects of home dampness and molds among Canadian children. *Am J Epidemiol* 1991;134:196-203.
32. Dales RE, Burnett R, Zwanenburg H. Adverse health effects among adults exposed to home dampness and molds. *Am Rev Respir Dis* 1991;143:505-9.
33. Kanhongkittiphon W, Mendell MJ, Gaffin JM, Wang G, Phipatanakul W. Indoor environmental exposures and exacerbation of asthma: an update to the 2000 review by the Institute of Medicine. *Environ Health Perspect* 2015;123:6-20.
34. Gent JF, Kezik JM, Hill ME, Tsai E, Li DW, Leaderer BP. Household mold and dust allergens: exposure, sensitization and childhood asthma morbidity. *Environ Res* 2012;118:86-93.
35. Pongracic JA, O'Connor GT, Muilenberg ML, Vaughn B, Gold DR, Kattan M, et al. Differential effects of outdoor versus indoor fungal spores on asthma morbidity in inner-city children. *J Allergy Clin Immunol* 2010;125:593-9.
36. Bundy KW, Gent JF, Beckett W, Bracken MB, Belanger K, Triche E, et al. Household airborne *Penicillium* associated with peak expiratory flow variability in asthmatic children. *Ann Allergy Asthma Immunol* 2009;103:26-30.
37. Black PN, Udy AA, Brodie SM. Sensitivity to fungal allergens is a risk factor for life-threatening asthma. *Allergy* 2000;55:501-4.
38. O'Driscoll BR, Hopkinson LC, Denning DW. Mold sensitization is common amongst patients with severe asthma requiring multiple hospital admissions. *BMC Pulm Med* 2005;5:4.
39. Knutsen AP, Bush RK, Demain JG, Denning DW, Dixit A, Fairs A, et al. Fungi and allergic lower respiratory tract diseases. *J Allergy Clin Immunol* 2012;129:280-91. quiz 92-3.
40. Neas LM, Dockery DW, Burge H, Koutrakis P, Speizer FE. Fungus spores, air pollutants, and other determinants of peak expiratory flow rate in children. *Am J Epidemiol* 1996;143:797-807.
41. Suphioglu C. Thunderstorm asthma due to grass pollen. *Int Arch Allergy Immunol* 1998;116:253-60.
42. Taylor PE, Flagan RC, Valenta R, Glovsky MM. Release of allergens as respirable aerosols: a link between grass pollen and asthma. *J Allergy Clin Immunol* 2002;109:51-6.
43. Taylor PE, Jonsson H. Thunderstorm asthma. *Curr Allergy Asthma Rep* 2004;4:409-13.
44. Taylor PE, Flagan RC, Miguel AG, Valenta R, Glovsky MM. Birch pollen rupture and the release of aerosols of respirable allergens. *Clin Exp Allergy* 2004;34:1591-6.
45. Grote M, Vrtala S, Niederberger V, Valenta R, Reichelt R. Expulsion of allergen-containing materials from hydrated rye grass (*Lolium perenne*) pollen revealed by using immunogold field emission scanning and transmission electron microscopy. *J Allergy Clin Immunol* 2000;105:1140-5.
46. Dannemiller KC, Gent JF, Leaderer BP, Peccia J. Influence of housing characteristics on bacterial and fungal communities in homes of asthmatic children [published online ahead of print March 31, 2015]. *Indoor Air*.
47. Wu AC, Lasky-Su J, Rogers CA, Klandermans BJ, Litonjua AA. Fungal exposure modulates the effect of polymorphisms of chitinases on emergency department visits and hospitalizations. *Am J Respir Crit Care Med* 2010;182:884-9.
48. Inal A, Karakoc GB, Altintas DU, Guvenmez HK, Aka Y, Geliskan R, et al. Effect of indoor mold concentrations on daily symptom severity of children with asthma and/or rhinitis monosensitized to molds. *J Asthma* 2007;44:543-6.
49. Turyk M, Curtis L, Scheff P, Contreras A, Coover L, Hernandez E, et al. Environmental allergens and asthma morbidity in low-income children. *J Asthma* 2006;43:453-7.
50. Tiesler CM, Thiering E, Tischer C, Lehmann I, Schaaf B, von Berg A, Heinrich J. Exposure to visible mould or dampness at home and sleep problems in children: Results from the LISAPlus study. *Environ Res* 2015;137:357-63.
51. Weinmayr G, Gehring U, Genuneit J, Büchele G, Kleiner A, Siebers R, et al. Dampness and moulds in relation to respiratory and allergic symptoms in children: results from Phase Two of the International Study of Asthma and Allergies in Childhood (ISAAC Phase Two). *Clin Exp Allergy* 2013;43:762-74.
52. Hagmolen of Ten Have W, van den Berg NJ, van der Palen J, van Aalderen WM, Bindels PJ. Residential exposure to mould and dampness is associated with adverse respiratory health. *Clin Exp Allergy* 2007;37:1827-32.
53. Kercsmar CM, Dearborn DG, Schluchter M, Xue L, Kirchner HL, Sobolewski J, et al. Reduction in asthma morbidity in children as a result of home remediation aimed at moisture sources. *Environ Health Perspect* 2006;114:1574-80.
54. Bernstein JA, Bobbitt RC, Levin L, Floyd R, Crandall MS, Shalwitz RA, et al. Health effects of ultraviolet irradiation in asthmatic children's homes. *J Asthma* 2006;43:255-62.
55. Bonner S, Matte TD, Fagan J, Andreopoulos E, Evans D. Self-reported moisture or mildew in the homes of Head Start children with asthma is associated with greater asthma morbidity. *J Urban Health* 2006;83:129-37.
56. Teach SJ, Crain EF, Quint DM, Hylan ML, Joseph JG. Indoor environmental exposures among children with asthma seen in an urban emergency department. *Pediatrics* 2006;117:S152-8.
57. Venn AJ, Cooper M, Antoniak M, Laughlin C, Britton J, Lewis SA. Effects of volatile organic compounds, damp, and other environmental exposures in the home on wheezing illness in children. *Thorax* 2003;58:955-60.
58. Jaakkola MS, Quansah R, Hugg TT, Heikkinen SA, Jaakkola JJ. Association of indoor dampness and molds with rhinitis risk: a systematic review and meta-analysis. *J Allergy Clin Immunol* 2013;132:1099-1110.e18.
59. Wen XJ, Balluz L, Mokdad A. Do obese adults have a higher risk of asthma attack when exposed to indoor mold? A study based on the 2005 Behavioral Risk Factor Surveillance System. *Public Health Rep* 2009;124:436-41.
60. Williamson IJ, Martin CJ, McGill G, Monie RD, Fennerty AG. Damp housing and asthma: a case-control study. *Thorax* 1997;52:229-34.

61. Pettigrew HD, Selmi CF, Teuber SS, Gershwin ME. Mold and human health: separating the wheat from the chaff. *Clin Rev Allergy Immunol* 2010;38:148-55.
62. Eickmeier O, Rieber N, Eckrich J, Hector A, Graeppler-Mainka U, Hartl D. Immune response, diagnosis and treatment of allergic bronchopulmonary aspergillosis in cystic fibrosis lung disease. *Curr Pharm Des* 2013;19:3669-78.
63. Patterson K, Streck ME. Allergic bronchopulmonary aspergillosis. *Proc Am Thorac Soc* 2010;7:237-44.
64. Chowdhary A, Agarwal K, Kathuria S, Gaur SN, Randhawa HS, Meis JF. Allergic bronchopulmonary mycosis due to fungi other than *Aspergillus*: a global overview. *Crit Rev Microbiol* 2014;40:30-48.
65. Gupta RK, Chandr A, Gautam PB. Allergic bronchopulmonary aspergillosis—a clinical review. *J Assoc Physicians India* 2012;60:46-51.
66. Mahdavinia M, Grammer LC. Management of allergic bronchopulmonary aspergillosis: a review and update. *Ther Adv Respir Dis* 2012;6:173-87.
67. Stewart AE, Hunsaker DH. Fungus-specific IgG and IgE in allergic fungal rhinosinusitis. *Otolaryngol Head Neck Surg* 2002;127:324-32.
68. Hamilos DL. Allergic fungal rhinitis and rhinosinusitis. *Proc Am Thorac Soc* 2010;7:245-52.
69. Fokkens WJ, Ebbens F, van Drunen CM. Fungus: a role in pathophysiology of chronic rhinosinusitis, disease modifier, a treatment target, or no role at all? *Immunol Allergy Clin North Am* 2009;29:677-88.
70. Thompson GR III, Patterson TF. Fungal disease of the nose and paranasal sinuses. *J Allergy Clin Immunol* 2012;129:321-6.
71. Schubert MS. Medical treatment of allergic fungal sinusitis. *Ann Allergy Asthma Immunol* 2000;85:90-7. quiz 7-101.
72. Thorp BD, McKinney KA, Rose AS, Ebert CS Jr. Allergic fungal sinusitis in children. *Otolaryngol Clin North Am* 2012;45:631-42. viii.
73. Greenberger PA. Mold-induced hypersensitivity pneumonitis. *Allergy Asthma Proc* 2004;25:219-23.
74. doPico GA, Reddan WG, Chmelik F, Peters ME, Reed CE, Rankin J. The value of precipitating antibodies in screening for hypersensitivity pneumonitis. *Am Rev Respir Dis* 1976;113:451-5.
75. Selman M, Lacasse Y, Pardo A, Cormier Y. Hypersensitivity pneumonitis caused by fungi. *Proc Am Thorac Soc* 2010;7:229-36.
76. Griesse M, Haug M, Hartl D, Teusch V, Glockner-Pagel J, Brasch F. Hypersensitivity pneumonitis: lessons for diagnosis and treatment of a rare entity in children. *Orphanet J Rare Dis* 2013;8:121.
77. Fracchia MS, El Saleeby CM, Murali MR, Sagar P, Mino-Kenudson M. Case records of the Massachusetts General Hospital. Case 9-2013. A 9-year-old boy with fever, cough, respiratory distress, and chest pain. *N Engl J Med* 2013;368:1141-50.
78. Watanuki Z, Okada S, Chiba S, Kamei K, Suzuki Y, Yamada N. Increased prevalence of high anti-*Cladosporium* antibody titers in interstitial lung diseases. *Tohoku J Exp Med* 2012;226:287-91.
79. Blatman KH, Grammer LC. Chapter 19: hypersensitivity pneumonitis. *Allergy Asthma Proc* 2012;33:S64-6.
80. Girard M, Cormier Y. Hypersensitivity pneumonitis. *Curr Opin Allergy Clin Immunol* 2010;10:99-103.